

<<双曲问题 (第2卷)>>

图书基本信息

书名：<<双曲问题 (第2卷)>>

13位ISBN编号：9787040345360

10位ISBN编号：7040345366

出版时间：2012-7

出版时间：高等教育出版社

作者：李大潜，江松 主编

页数：759

字数：520000

版权说明：本站所提供下载的PDF图书仅提供预览和简介，请支持正版图书。

更多资源请访问：<http://www.tushu007.com>

## &lt;&lt;双曲问题 (第2卷)&gt;&gt;

## 内容概要

This two-volume book is devoted to mathematical theory, numerics and applications of hyperbolic problems. Hyperbolic problems have not only a long history but also extremely rich physical background. The development is highly stimulated by their applications to Physics, Biology, and Engineering Sciences; in particular, by the design of effective numerical algorithms. Due to recent rapid development of computers, more and more scientists use hyperbolic partial differential equations and related evolutionary equations as basic tools when proposing new mathematical models of various phenomena and related numerical algorithms.

This book contains 80 original research and review papers which are written by leading researchers and promising young scientists, which cover a diverse range of multidisciplinary topics addressing theoretical, modeling and computational issues arising under the umbrella of "Hyperbolic Partial Differential Equations". It is aimed at mathematicians, researchers in applied sciences and graduate students.

## &lt;&lt;双曲问题 (第2卷)&gt;&gt;

## 书籍目录

Volume 2

Contributed Talks

Alin, a Chertock, Jian, -Guo Liu, Terran, ce Pendleton

Convergence Analysis of the Particle Method for the  
Camassa-Holm Equation

Paolo Corti

Stable Numerical Scheme for the Magnetic Induction Equation  
with Hall Effect

Andreas Dedner, Christoph Gersbacher

A Local Discontinuous Galerkin Discretization for Hydrostatic  
Free Surface Flows Using  $\alpha$ -transformed Coordinates

Don, atella Don, atelli, Pierangelo Marcati

Analysis of Quasineutral Limits

Renjun Duan, Robert M. Strain

On the Full Dissipative Property of the Vlasov-Poisson-Boltzmann  
System

P. Engel, C. Rohde

On the Space-Time Expansion Discontinuous Galerkin  
Method

Ulrik Skre Fjordholm

Energy Conservative and Stable Schemes for the Two-layer  
Shallow Water Equations

Jan, Giesselmann

Sharp Interface Limits for Korteweg Models

Nan Jiang

On the Convergence of Semi-discrete High Resolution Schemes  
with Superbee Flux Limiter for Conservation Laws

Song Jiang, Qian, gchang Ju, Fucai

Asymptotic Limits of the Compressible Magnetohydrodynamic  
Equations

Chunhua Jin, Tong Ya'ng, Jin, gxue Yitn,

Waiting Time for a Non-Newtonian Polytopic Filtration  
Equation with ConvectionQuansen Jiu, Yi Wang, Zhouping Xin, Stability of Rarefaction Waves to  
the 1D Compressible Navier-Stokes Equations with Density-dependent  
Viscosity

E. A. John, son, J. A. Rossmann

Ten-moment Two-fluid Plasma Model Agrees Well with  
PIC/Vlasov in GEM Problem

Frederike Kissling, Christian Rohde

Numerical Simulation of Nonclassical Shock Waves in Porous  
Media with a Heterogeneous Multiscale Method

Ujjwal Koley

Implicit Finite Difference Scheme for the Magnetic  
Induction Equation

<<双曲问题 (第2卷)>>

Anastasia Korshun,ova,Olga Rozanova

On Effects of Stochastic Regularization for the Pressureless

Gas Dynamics

Dietmar Kroener

Jump Conditions across Phase Boundaries for the Navier-Stokes

Korteweg system

Kai Krycki,Martitn, Frank

Numerical Treatment of a Non-classical Transport Equation

Modelling Radiative Transfer in Atmospheric Clouds

.....

Volume 1

## 章节摘录

版权页：插图： The resolution of the rarefaction wave is more troublesome, because the rarefaction advances into a constant state, where the mesh is very coarse: the mesh should be refined fast enough to resolve the corner of the rarefaction correctly. Again, the AdLim strategy performs very poorly, and the CT scheme produces a step in the middle of the rarefaction. Again, ENO and S&E ( not shown ) are very similar to MM. On the other hand, the rarefaction wave is well resolved by the uniform grid ( see also the very low values of the entropy production, which is plotted on the lower part of the figure ) . Obviously, the cost of the solution with the uniform grid is much higher, since it is using a much larger number of grid points. ( Note that for  $S_{ref}=0.001$ , the rarefaction wave is correctly resolved by all schemes and allowing adaptive schemes to use cells of width  $1/8192$  as in the reference solution would improve dramatically the resolution of all schemes. ) Figure 3.2 shows the complete density component of the solution for the shock-acoustic wave interaction ( a ) together with the cell levels employed. Note that in the area behind the big shock, cells are kept to the maximum refinement level, while some coarsening is allowed in the smooth area between the small shocks on the left. On the right part of the figure ( b ) , we show the number of cells used by the different adaptive algorithms as a function of time. It is clear that all algorithms use approximately the same number of cells, and this number increases with time, as the solution evolves and more structures emerge. In Figure 3.3 we enlarge two areas of the solution to highlight the differences among the schemes. We observe that the MM method performs quite poorly on this problem, because of the clipping of extrema by the MinMod limiter: the acoustic wave is smeared due to grid coarsening before entering into the shock and the waves do not have the correct amplitude after the shock ( b ) .

<<双曲问题 (第2卷)>>

编辑推荐

《双曲问题:理论、数值方法及应用(第2卷)》由高等教育出版社出版。

<<双曲问题 (第2卷)>>

版权说明

本站所提供下载的PDF图书仅提供预览和简介, 请支持正版图书。

更多资源请访问:<http://www.tushu007.com>