

<<啤酒大麦品质的遗传和改良>>

图书基本信息

书名：<<啤酒大麦品质的遗传和改良>>

13位ISBN编号：9787308063821

10位ISBN编号：7308063828

出版时间：2009-6

出版时间：浙江大学出版社

作者：张国平，李承道 主编

页数：296

字数：660000

版权说明：本站所提供下载的PDF图书仅提供预览和简介，请支持正版图书。

更多资源请访问：<http://www.tushu007.com>

<<啤酒大麦品质的遗传和改良>>

内容概要

Genetics and Improvement of Barley Malt Quality presents up-to-date developments in barley production and breeding . The book is divided into nine chapters , including barley production and consumption , germplasm and utilization , chemical composition , protein and protein components , carbohydrates and sugars , starch degrading enzymes , endosperm cell walls and malting quality , genomics and malting quality improvement , and marker-assisted selection for malting quality The information will be especially useful to barley breeders , malsters , brewers , biochemists , barley quality specialists , molecular geneticists , and biotechnologists . This book may also serve as reference text for post-graduate students and barley researchers . The authors for each chapter are the experts and frontier researchers in the specific areas .

<<啤酒大麦品质的遗传和改良>>

作者简介

Professor Guoping Zhang , is a barley breeder and crop physiologist in Department of Agronomy , Zhejiang University of China . Dr . Chengdao Li is a senior molecular geneticist and barley breeder in Department of Agriculture & Food , Western Australia . He is also an adjunct professor in Murdoch university of Australia and Zhejiang University of China .

<<啤酒大麦品质的遗传和改良>>

书籍目录

1 Barley Production and Consumption 1.1 Introduction 1.2 World Barley Production 1.3 World Barley Trade 1.4 World Barley Consumption 1.4.1 Feed Consumption 1.4.2 Malting Barley Consumption 1.4.3 Food Consumption 1.4.4 Other Industrial Uses References 2 Barley Germplasm and Utilization 2.1 Origin and Evolution of Cultivated Barley 2.1.1 The Taxonomy of Barley 2.1.2 Origin of the Cultivated Barley 2.1.3 Evolution and Domestication of Cultivated Barley 2.2 Annual Wild Barley 2.2.1 Annual Wild Barley in the Near East Fertile Crescent 2.2.2 The Annual Wild Barley in Qing-Tibetan Plateau 2.2.3 Other Annual Wild Barley Species 2.2.4 Prospects 2.3 Perennial Wild Barley Germplasm 2.3.1 The Variation of Target Traits 2.3.2 Approaches for Utilization of Perennial Wild Barleys 2.4 Cultivated Barleys 2.4.1 Primitive Barleys or Landraces 2.4.2 Commercial Varieties 2.4.3 Mutants References 3 Chemical Composition in Barley Grains and Malt Quality 3.1 Introduction 3.2 Physical Structure 3.2.1 Grain Size 3.2.2 Dormancy 3.2.3 Grain Hardness 3.3 The Internal Structure 3.3.1 The Embryo 3.3.2 The Aleurone 3.3.3 The Barley Husk 3.4 The Barley Endosperm 3.4.1 Barley Carbohydrates 3.4.2 Grain Protein 3.5 Malt Quality 3.5.1 Diastatic Power 3.5.2 α -amylase 3.5.3 β -amylase 3.5.4 Limit Dextrinase 3.5.5 α -glucosidase 3.5.6 β -glucanase 3.5.7 Proteinase 3.5.8 Hot Water Extract 3.6 Malted Barley Quality Traits 3.7 Conclusion References 4 Identification of Barley Varieties by Protein Profiling 4.1 Introduction 4.2 Extraction of Proteins for Protein Profiling 4.3 Characteristics of Barley Protein Profiles 4.3.1 Peak Size Calling 4.3.2 Location and Year 4.3.3 Phylogeny 4.4 Discussion 4.4.1 Specificity of Variety Identification 4.4.2 Location and Year 4.4.3 Malting versus Malt 4.4.4 Peak Inheritance 4.4.5 Contamination 4.4.6 Automation 4.5 Conclusion References 5 α -glucans and Arabinoxylans 6 The Properties and Genetics of Barley Malt Starch 7 The Role of Endosperm Cell Walls in Barley Malting Quality 8 Barley Genomics and Malting Quality Improvement 9 Genetic Improvement of Malting Quality through Conventional Breeding and Marker-assisted Selection Index

<<啤酒大麦品质的遗传和改良>>

章节摘录

插图：Barley has many economic uses today. Barley is produced primarily as animal feed. For example, over half of the barley is used for livestock feed in the United States. Barley as feed has the same nutritive value as corn. Barley is high in carbohydrates, with moderate amounts of protein, calcium and phosphorus. It also has small amounts of vitamin B. The entire barley kernel is used as feed after it has been steam rolled or gone through a grinding process. By products from the brewing process and malt sprouts are also used in livestock feed. The two-rowed barley is most often used for animal feed because it produces higher weight and superior kernel production. Barley is also used in the production of beer and some wines. About 25% of the cultivated barley in the United States is used for malting, with about 80% used in beer production, 14% used in distilled alcohol production, and 6% used for malt syrup, malted milk, and breakfast foods. A small amount of the barley is used for human food in the form of pearl barley or flour for porridge. Sometimes, barley is grown as a hay crop in some areas. Only the smooth varieties and awnless varieties are used in hay production. Winter barley also can be used for hay if pasteurized before the stems start to elongate. The amount of barley used for ethanol has been increased significantly in the last decade, especially in some EU countries (www.biofuelreview.com).

<<啤酒大麦品质的遗传和改良>>

编辑推荐

《啤酒大麦品质的遗传和改良(英文版)(精)》由浙江大学出版社出版。

<<啤酒大麦品质的遗传和改良>>

版权说明

本站所提供下载的PDF图书仅提供预览和简介，请支持正版图书。

更多资源请访问:<http://www.tushu007.com>