

<<中国油气区第三系>>

图书基本信息

书名：<<中国油气区第三系>>

13位ISBN编号：9787502116699

10位ISBN编号：7502116699

出版时间：1997-05-01

出版时间：叶得泉、Zhong Xiaochun、Yao Yimin、Yang Fan 石油工业出版社（1997-05出版）

作者：叶得泉 等著

版权说明：本站所提供下载的PDF图书仅提供预览和简介，请支持正版图书。

更多资源请访问：<http://www.tushu007.com>

<<中国油气区第三系>>

内容概要

《中国油气区第三系（英文版）》主要包括：CHARACTERISTICS OF THE DISTRIBUTION OF TERTIARY IN CHINA OIL-AND GAS-BEARING REGIONS、TERTIARY BIOTAS OF CHINA OIL-AND GAS-BEARING REGIONS、APPLICATION OF SEISMIC STRATIGRAPHY, MAGNETOSTRATIGRAPHY ISOTOPLC CHRONOSTRATIGRAPHY AND EVENT STRATIGRAPHY METHODS TO STRATIGRAPHIC CORRELATIONS、CORRELATIONS OF TERTIARY STRATA IN CHINA OIL-AND GAS-BEARING REGIONS TERTIARY PALAEOENVIRONMENT OF CHINA OIL-AND GAS-BEARING REGIONS等。

<<中国油气区第三系>>

书籍目录

CHAPTER 1 CHARACTERISTICS OF THE DISTRIBUTION OF TERTIARY IN CHINA OIL AND GAS-BEARING REGIONS 1.1 Widespread Depositional Basins 1.2 Dominance of Nonmarine Deposits, with a Variety of Deposition Types 1.3 Tertiary Development under Control of Tectonic Movement 1.4 Extremely Abundant Fossils 1.5 Frequent Volcanic Activities in East China with an Extensive Distribution of Volcanic Rocks 1.6 Rich Mineral Resources CHAPTER 2 TERTIARY BIOTAS OF CHINA OIL AND GAS-BEARING REGIONS 2.1 Trilobites 2.2 Foraminifers 2.3 Gastropods 2.4 Bivalves 2.5 Vertebrates 2.6 Sporopollen 2.7 Dinoflagellates and Acritarchs 2.8 Charophytes 2.9 Calcareous Nannofossils 2.10 Fossil Plants CHAPTER 3 APPLICATION OF SEISMIC STRATIGRAPHY, MAGNETOSTRATIGRAPHY, ISOTOPIC CHRONOSTRATIGRAPHY AND EVENT STRATIGRAPHY METHODS TO STRATIGRAPHIC CORRELATIONS 3.1 Application of Seismic Stratigraphy to Stratigraphic Correlations 3.2 Isotopic Geological Age Determination of Volcanic Rocks 3.3 Age-Determination of Palaeogeomagnetism 3.4 Application of Stable Isotope Stratigraphy and Event Stratigraphy CHAPTER 4 CORRELATIONS OF TERTIARY STRATA IN CHINA OIL AND GAS-BEARING REGIONS 4.1 Typical Tertiary Sequences 4.2 Basis for Tertiary Stratigraphical Correlations 4.3 Correlations of Tertiary Strata 4.4 Correlations with Tertiary Normal Sequence of Europe and America 4.5 Discussion on Tertiary Boundaries CHAPTER 5 TERTIARY PALAEOENVIRONMENT OF CHINA OIL AND GAS-BEARING REGIONS 5.1 Tertiary Palaeovegetation and Palaeoclimate 5.2 Tertiary Palaeobiogeographical Provinces and their Controlling Factors in China 5.3 Discussions on the Paleogene "Transgression" in East China Oil and Gas-bearing Regions CHAPTER 6 DIVISION OF TERTIARY OIL-GENERATING PERIODS AND PREDICTION OF OIL AND GAS PROSPECTS IN OIL AND GAS-BEARING REGIONS OF CHINA 6.1 Tertiary Oil Source Rocks Distribution in Chinese Oil and Gas-bearing Regions 6.2 Influential Factors on Development of Dark-Coloured Strata and Hydrocarbon Potential 6.3 Division of Tertiary Oil-Generating Periods 6.4 Application of Stratigraphical Palaeontological Studies to Prediction of Oil and Gas Potential 6.5 Prediction of Tertiary Oil and Gas Prospects CONCLUDING REMARKS REFERENCES

章节摘录

版权页： 插图： 1.1.7 Northern South China Sea Shelf Oil and Gas-bearing Region This region mainly includes the Zhujiangkou (Pearl River Mouth), Beibuwan, Yinggehai and Southeastern Qiongya Basins in the Northern sea area of the South China Sea together with some other local areas of Hainan and Guangdong Provinces. In this region the Paleogene mainly belongs to continental deposits; the transgression began from Late Oligocene and became further expanded up to the Neogene. As a result, in this region the Neogene is mainly developed with marine deposits, which are extensively distributed, with a thickness of more than 3,500m. This region also represents a distributive area of variegated elastic rocks. The coast-line of Mainland China is as long as 18,000km, with an extremely vast expanse of territorial and neighbouring sea areas. Within the sea areas of Bohai, Huanghai, East China Sea and South China Sea, there are 16 depositional basins with developed Tertiary, mostly belonging to those of inheritance type and newborn type, with a total area attaining 1,955,500km². It is doubtless, therefore, that there exist extremely broad prospects in the Tertiary oil and gas exploration over the sea areas of China. However, in the Xizang Region, the Tertiary is not very well-developed, and therefore exploration for oil and gas has not been carried out to a great extent therein. Accordingly, the Xizang Region is not included in the research Project on the Tertiary of China Oil and Gas-bearing Regions. In the Xizang Region, there are 18 Mesozoic and Cenozoic depositional basins, including 9 Tertiary basins of the newborn type, which are generally studied in a lower degree. 1.2 Dominance of Nonmarine Deposits, with a Variety of Deposition Types In China, the nonmarine Tertiary, with a great thickness, is extensively distributed all over the depositional basins of various genetic types throughout the country. As compared with marine deposits, the continental deposits obviously occupy a dominant position, while marine deposits are distributed within a relatively narrow scope.

<<中国油气区第三系>>

编辑推荐

《中国油气区第三系(英文版)》由石油工业出版社出版。

<<中国油气区第三系>>

版权说明

本站所提供下载的PDF图书仅提供预览和简介，请支持正版图书。

更多资源请访问:<http://www.tushu007.com>